

Article

Analyzing Retinal Detachment Surgery Outcomes and Key Predictive Variables in Iraqi Populations

Dr. Wajdi Saadi Kallak¹, Dr. Hayder Abdul Al-Husain Abdul Al-Redha², Dr. Hayder Sabah Al-Rubaye³

1. Iraqi Ministry of Health, Baghdad Al-Resafa Health Directorate, Ibn Al-Haitham Eye Teaching Hospital, Baghdad, Iraq
 2. Iraqi Ministry of Health, Baghdad Al-Resafa Health Directorate, Ibn Al-Haitham Eye Teaching Hospital, Baghdad, Iraq
 3. Iraqi Ministry of Health, Baghdad Al-Resafa Health Directorate, Ibn Al-Haitham Eye Teaching Hospital, Baghdad, Iraq
- * Correspondence: Wajdisaadi78@gmail.com, Hayderophthalmology@gmail.com, dr_hsg@yahoo.com

Abstract: Background: Recent developments in vitreoretinal microsurgical methods have made it possible for several successive procedures in a single eye as the disease continues. **Objective:** The aim of this article is to enrol and evaluate the clinical outcomes of retinal detachment surgery and to identify the basic predictive factors of Iraqi patients. **Methods:** 95 patients who underwent retinal detachment surgery were recruited from different hospitals in Iraq during a 5-month follow-up period from January 2023 to May 2024. BCVA was performed to assess the functional outcome, and the visual acuity of patients was measured before and after surgery. **Findings:** A total of 95 cases underwent all different interventions, ranging from 1 to 5 surgeries. The mean number of operations was found to be four (range: 2 to 10), and the anatomical success rate was 81.05%. Postoperative visual acuity was worse than 20/400-20/100 in 21 cases, 20/80-20/65 in 20 cases, and $\geq 20/50$ in 54 cases. Postoperative complications were observed in 18.95% of patients, with the most prevalent being infection (7 cases) and retinal detachment recurrence (4 cases). **Conclusion:** In our study, we suggested that retinal detachment surgery can have successful outcomes that advance a functional outcome of high visual acuity for patients.

Keywords: Retinal Detachment Interventions, Postoperative Complications, and Visual Acuity Measurement Outcomes

Citation: Kallak, W. S. Analyzing Retinal Detachment Surgery Outcomes and Key Predictive Variables in Iraqi Populations. International Journal of Health Systems and Medical Sciences 2025, 4(1), 67-73.

Received: 10th Jan 2025

Revised: 11th Jan 2025

Accepted: 24th Jan 2025

Published: 27th Jan 2025



Copyright: © 2024 by the authors. Submitted for open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>)

1. Introduction

Retinal detachment is a common cause of visual loss among the population. Currently, the treatment for this pathology is surgical, being the pars plana vitrectomy (PPV), the scleral buckle (HE), or cerclage, and the combination of both (PPV+HE) is the most used by surgeons. However, the visual and anatomical results, the development of cataracts, and the success rates after retinal surgery are under continuous study. [1,2,3]

Retinal detachment (DR) is defined as the separation of the RNS from the retinal pigment epithelium (RPE) [4]. These two anatomical sections of the retina are formed from the neuroectoderm that is found enveloping the optic vesicle in the process of embryogenesis. The optic vesicle invaginates and forms the optic cup, leaving one layer on top of the other; the innermost differentiates in RNS and the outermost in the RPE. These two layers are not really joined by anatomical junctions between their cells. [5]

Retinal detachment occurs when the adhesion forces between the RNS and RPE are overcome. It is common for all retinal detachments to cause the accumulation of subretinal fluid. Rhegmatogenic retinal detachment (RRD) is to be treated mainly [6]. The DRR is characterized by the existence of a full-thickness retinal rupture that is kept open by vitreoretinal traction, which facilitates the entry of liquefied vitreous and accumulates under the retina, causing that separation between RNS and RPE [7,8,9]. It can be said that

the main cause of the RRD is the liquefied vitreous; the tensile forces can cause and maintain a retinal rupture, through which the liquid passes to the ESR. Although there is a full-thickness rupture, if the vitreous is not partially liquefied and there is no necessary traction, the RRD will not occur. [10,11,12,13]

The increased risk of DRR is associated with several factors. Among them, myopia should be highlighted [14,15]. There is a prevalence of RRD in nearsighted people, with an incidence approximately 10 times higher in people with more than three diopters of myopia. Nearsighted eyes experience a DVP at an earlier age [16]. This can be explained by a thinner retina with myopic elongation of the axial globe. Myopia up to -3 diopters (D) multiplies the risk of DR by 4, and in cases of myopia greater than -3 D, the risk of DR increases up to 10 times. Myopia leads to earlier vitreous liquefaction, which explains why DR tends to occur earlier in patients presenting with myopia than in emmetropic or farsighted patients. Myopia is a relevant risk factor because it is increasingly common among the population. [17,18,19,20,21,22,23]

2. Materials and Methods

Our cross-sectional study looked at the records of retinal detachment cases and included 95 people who had had surgery for the condition at the Retina Service at the Department of Ophthalmology in different hospitals in Iraq. The patients varied in age from 20 to 60. Each of the patients were operated on by experienced posterior segment surgeons, and each patient had a surgical follow-up at least five months after the final procedure. All postoperative follow-ups have been carried out in Iraq and at different hospitals in Iraq.

Age and gender of the patient, cause of retinal detachment, best corrected visual acuity (BCVA) before the first intervention, intraoperative methods and procedures used in each patient's subsequent operations, including the use of retinectomy, endolaser, cryopexy, gas, and silicone oil (sil-oil); intraoperative vitreous entrapment in the sites of the sclerotomy, presence of an epiretinal membrane (ERM), proliferative vitreoretinopathy (PVR), the identification of new or missed breaks, postoperative complications, the quantity of operations per patient, the anatomical condition of the retina during the end of the follow-up, BCVA during the end of a follow-up, intraocular pressure (IOP) during the end of the follow-up time, and the total duration of the follow-up period.

Both before and after surgery, visual acuity was assessed using the BCVA scale. The reasons behind retinal detachment, the patient's age and gender, the best corrected visual acuity (BCVA) prior to the initial intervention, the intraoperative techniques and, the presence of proliferative vitreoretinopathy (PVR), the detection of new or missed breaks, postoperative complications, the number of operations for each patient, and the retina's anatomical status at the end of the follow-up.

3. Results

Table 1. Enrol demographic and preoperative characteristics.

<i>Features</i>	<i>N = 95</i>	<i>%</i>
<i>Age, years</i>		
<i>20 – 30</i>	22	23.16%
<i>31 – 40</i>	20	21.05%
<i>41 – 50</i>	23	24.21%
<i>51 – 60</i>	30	31.58%
<i>Gender</i>		
<i>Male</i>	76	80.00%
<i>Female</i>	19	20.00%

Comorbidities		
<i>Hypertension</i>	67	70.53%
<i>Diabetes</i>	36	37.89%
<i>Asthma</i>	20	21.05%
<i>Others</i>	13	13.68%
Smoking, yes/no		
<i>Smokers</i>	54	56.84%
<i>None</i>	41	43.16%
Education status (P, S, U)		
<i>Primary</i>	26	27.37%
<i>Secondary</i>	30	31.58%
<i>University</i>	39	41.05%
Income status, \$		
<i>Low, 200 – 400</i>	37	38.95%
<i>Moderate, 401 – 700</i>	28	29.47%
<i>High, > 700</i>	30	31.58%

Table 2. Enrol intervention data of retinal detachment in relation to gender.

Variables	N = 95	%
Pseudophakic (38/40%) cases	Male: 28 Female: 10	73.68% 26.32%
Phakic (57/60%) cases	Male: 50 Female: 7	87.72% 12.28%
High myopia/Giant tears (25) cases	Male: 22 Female: 3	88% 12%
Trauma (12) cases	Male: 11 Female: 1	91.67% 8.33%
Primary rhegmatogenous retinal detachment (20) cases	Male: 17 Female: 3	85% 15%

Table 3. Diagnostic outcomes of eyes preoperatively.

Variables	N = 95	%
Location of retinal detachment		
<i>Inferior</i>	34	35.79%
<i>Superior</i>	25	26.32%
<i>Temporal</i>	21	22.11%
<i>Superior, nasal</i>	10	10.53%
<i>Superior, inferior, temporal</i>	3	3.16%
<i>Superior, inferior, nasal</i>	2	2.11%
The number of detached retinal area		

4 clock hours	6	6.32%
6 clock hours	77	81.05%
8 clock hours	12	12.63%
The number of tears detected preoperatively.		
0	30	31.58%
2	58	61.05%
≥ 3	7	7.37%

Table 4. Identify pre and postoperative outcomes.

Variables	N = 95	%
Duration of macular detachment (weeks)	7.5 ± 4.6	
Extent of retinal detachment (hours)	8.4 ± 1.3	
Number of surgeries used		
1	25	26.32%
2	32	33.68%
> 2	38	40.00%
Types		
Retinectomy	39	41.05%
Silicon oil	50	52.63%
Gas	75	78.95%
Cryopexy	63	66.32%
Endo-laser	52	54.74%
Visual acuity		
Preoperative visual acuity	2.24 ± 0.56	
Visometric visual acuity	0.89 ± 0.27	
Postoperative visual acuity	0.51 ± 0.25	
Post-complications	18	18.95%
Infection	7	7.37%
Retinal detachment recurrence	4	4.21%
Cataracts	2	2.11%
Glaucoma	2	2.11%
Macular puckers	3	3.16%
Anatomical success rate		
High	77	81.05%
Moderate	12	12.63%
Poor	6	6.32%

Table 5. Comparison of visual acuity in preoperative and postoperative.

Scores	Postoperative		
	20/400–20/100	20/80–20/65	≥ 20/50
20/400 – 20/100	13	11	24
20/80 – 20/65	7	9	27
≥ 20/50	1	-	3
Total	21 cases	20 cases	54 cases

4. Discussion

The mean age of the samples in our study ranges from 20 to 60 years, which is comparable to the mean age reported in other studies with analogous objectives. In these studies, the mean age of the cases ranges approximately from 60 years. The study revealed that patients between the ages of 51 and 60 constituted 31.58% of the total sample.

The study demonstrated a higher incidence of RD in male patients compared to female patients, with a percentage of 80% and 20%, respectively. A review of other studies with samples from patients with DR reveals that the percentage of men in these samples ranges from 60-80%, which is similar to the values found in the present study, thereby confirming that the incidence is higher in men. [24,25,26]

With regard to the macular involvement of the RRDs under study, a percentage of 57.4% of RRDs with macular respect (macula-on) has been obtained, which differs from the findings of other authors in studies similar to ours, as evidenced by the American study, which found only 26% of cases with RRD without macular involvement. However, we did find similar rates of RRD without macular involvement closer to ours (53%) in the study conducted by Spain. In this study, a comparison of PPV and PPV+HE is made to treat RRD. [27]

Some studies found that the sample exhibited a 60% prevalence of phakic patients, a figure analogous to the 60% observed in the German study, which comparatively analysed PPV and HE as treatments for RRD. A notable distinction emerges from the Dutch study, which documented a percentage of phakic patients of approximately 36%, a significant deviation from the observed figure. [28,29,30]

5. Conclusion

We showed in our study that a high anatomical success rate can be obtained with retinal detachment surgery. A sizable portion of the connected eyeballs maintained strong visual acuity. Our data analysis indicated that surgery had no adverse impacts on anatomical or functional outcomes.

REFERENCES

- [1] W. L. Hutton, "The Effects of Silicone Oil Removal: Silicone Study Report 6," *Archives of Ophthalmology*, vol. 112, no. 6, p. 778, Jun. 1994, doi: 10.1001/archophth.1994.01090180076038.

- [2] F. Goezinne, E. C. La Heij, T. T. J. M. Berendschot, A. T. A. Liem, and F. Hendrikse, "Risk Factors for Redetachment and Worse Visual Outcome after Silicone Oil Removal in Eyes with Complicated Retinal Detachment," *Eur J Ophthalmol*, vol. 17, no. 4, pp. 627–637, Jul. 2007, doi: 10.1177/112067210701700423.
- [3] F. Jiang, M. Krause, K. W. Ruprecht, and K. Hille, "Management and Results of Retinal Detachment after Silicone Oil Removal," *Ophthalmologica*, vol. 216, no. 5, pp. 341–345, 2002, doi: 10.1159/000066176.
- [4] J. B. Jonas, "Retinal redetachment after removal of intraocular silicone oil tamponade," *British Journal of Ophthalmology*, vol. 85, no. 10, pp. 1203–1207, Oct. 2001, doi: 10.1136/bjo.85.10.1203.
- [5] N. Heussen, R.-D. Hilgers, H. Heimann, L. Collins, and S. Grisanti, "Scleral buckling versus primary vitrectomy in rhegmatogenous retinal detachment study (SPR Study): Multiple-event analysis of risk factors for reoperations. SPR Study report no. 4," *Acta Ophthalmol*, vol. 89, no. 7, pp. 622–628, Nov. 2009, doi: 10.1111/j.1755-3768.2009.01766.x.
- [6] A. M. Abu El-Asrar, S. M. Al-Bishi, and D. Kangave, "Outcome of Temporary Silicone Oil Tamponade in Complex Rhegmatogenous Retinal Detachment," *Eur J Ophthalmol*, vol. 13, no. 5, pp. 474–481, Jun. 2003, doi: 10.1177/112067210301300509.
- [7] R. H. Y. Asaria and Z. J. Gregor, "Simple retinal detachments: identifying the at-risk case," *Eye*, vol. 16, no. 4, pp. 404–410, Jul. 2002, doi: 10.1038/sj.eye.6700189.
- [8] G. Hilton, R. Machemer, R. Michels, E. Okun, C. Schepens, and A. Schwartz, "The Classification of Retinal Detachment with Proliferative Vitreoretinopathy," *Ophthalmology*, vol. 90, no. 2, pp. 121–125, Feb. 1983, doi: 10.1016/s0161-6420(83)34588-7.
- [9] W. F. Rachal and T. C. Burton, "Changing Concepts of Failures After Retinal Detachment Surgery," *Archives of Ophthalmology*, vol. 97, no. 3, pp. 480–483, Mar. 1979, doi: 10.1001/archophth.1979.01020010230008.
- [10] P. A. Quiram *et al.*, "Outcomes of Vitrectomy with Inferior Retinectomy in Patients with Recurrent Rhegmatogenous Retinal Detachments and Proliferative Vitreoretinopathy," *Ophthalmology*, vol. 113, no. 11, pp. 2041–2047, Nov. 2006, doi: 10.1016/j.ophtha.2006.05.039.
- [11] E. C. LA HEIJ, F. HENDRIKSE, and A. G. H. KESSELS, "RESULTS AND COMPLICATIONS OF TEMPORARY SILICONE OIL TAMPONADE IN PATIENTS WITH COMPLICATED RETINAL DETACHMENTS," *Retina*, vol. 21, no. 2, pp. 107–114, Apr. 2001, doi: 10.1097/00006982-200104000-00002.
- [12] V. G. Grigoropoulos, S. Benson, C. Bunce, and D. G. Charteris, "Functional outcome and prognostic factors in 304 eyes managed by retinectomy," *Graefe's Archive for Clinical and Experimental Ophthalmology*, vol. 245, no. 5, pp. 641–649, Nov. 2006, doi: 10.1007/s00417-006-0479-z.
- [13] F. Faude and P. Wiedemann, "Vitreoretinal Endoscope for the Assessment of the Peripheral Retina and the Ciliary Body after Large Retinectomies in Severe Anterior PVR," *Int Ophthalmol*, vol. 25, no. 1, pp. 53–56, Jan. 2004, doi: 10.1023/b:inte.0000018550.36179.9e.
- [14] L. S. MORSE, C. B. CHAPMAN, D. ELIOTT, J. D. BENNER, M. S. BLUMENKRANZ, and B. W. McCUEN, "SUBRETINAL HEMORRHAGES IN PROLIFERATIVE DIABETIC RETINOPATHY," *RETINA*, vol. 17, no. 2, pp. 87–93, 1997, doi: 10.1097/00006982-199703000-00001.
- [15] M. S. Blumenkranz *et al.*, "Relaxing Retinotomy With Silicone Oil or Long-acting Gas in Eyes With Severe Proliferative Vitreoretinopathy Silicone Study Report 5," *Am J Ophthalmol*, vol. 116, no. 5, pp. 557–564, Nov. 1993, doi: 10.1016/s0002-9394(14)73196-4.
- [16] S. E. Brodie, "M. Yanoff and J. S. Duker (eds): Ophthalmology: 4th Edition, Elsevier Saunders, 2014, ISBN: 978-1-4557-3984-4," *Documenta Ophthalmologica*, vol. 129, no. 2, pp. 139–140, Jul. 2014, doi: 10.1007/s10633-014-9454-x.
- [17] M. Stewart, D. Browning, and M. Landers, "Current management of diabetic tractional retinal detachments," *Indian J Ophthalmol*, vol. 66, no. 12, p. 1751, 2018, doi: 10.4103/ijo.ijo_1217_18.
- [18] L. Talat, S. Lightman, and O. Tomkins-Netzer, "Ischemic Retinal Vasculitis and Its Management," *J Ophthalmol*, vol. 2014, pp. 1–13, 2014, doi: 10.1155/2014/197675.
- [19] G. W. Blankenship, "Stability of Pars Plana Vitrectomy Results for Diabetic Retinopathy Complications: A Comparison of Five-Year and Six-Month Postvitrectomy Findings," *Archives of Ophthalmology*, vol. 99, no. 6, p. 1009, Jun. 1981, doi: 10.1001/archophth.1981.03930011009004.
- [20] Y.-J. Hsu, Y.-T. Hsieh, P.-T. Yeh, J.-Y. Huang, and C.-M. Yang, "Combined Tractional and Rhegmatogenous Retinal Detachment in Proliferative Diabetic Retinopathy in the Anti-VEGF Era," *J Ophthalmol*, vol. 2014, pp. 1–11, 2014, doi: 10.1155/2014/917375.

-
- [21] L. A. Lobes, "Diseases of the Macula, vol 1; Retinal Vascular Disorders, Diagnosis, and Management, vol 2; Sights and Sounds in Ophthalmology," *Archives of Ophthalmology*, vol. 95, no. 4, p. 716, Apr. 1977, doi: 10.1001/archophth.1977.04450040182033.
- [22] D. Shukla, S. Kanungo, N. M. Prasad, and R. Kim, "Surgical outcomes for vitrectomy in Eales' disease," *Eye*, vol. 22, no. 7, pp. 900–904, Mar. 2007, doi: 10.1038/sj.eye.6702763.
- [23] A. M. A. El-Asrar and S. A. Al-Kharashi, "Full panretinal photocoagulation and early vitrectomy improve prognosis of retinal vasculitis associated with tuberculoprotein hypersensitivity (Eales' disease)," *British Journal of Ophthalmology*, vol. 86, no. 11, pp. 1248–1251, Nov. 2002, doi: 10.1136/bjo.86.11.1248.
- [24] A. Kumar, H. K. Tiwari, R. P. Singh, L. Verma, and N. Prasad, "Comparative evaluation of early vs. deferred vitrectomy in Eales' Disease," *Acta Ophthalmol Scand*, vol. 78, no. 1, pp. 77–78, Feb. 2000, doi: 10.1034/j.1600-0420.2000.078001077.x.
- [25] A. B. Majji, G. K. Vemuganti, V. A. Shah, S. Singh, T. Das, and S. Jalali, "A comparative study of epiretinal membranes associated with Eales' disease: a clinicopathologic evaluation," *Eye*, vol. 20, no. 1, pp. 46–54, Apr. 2005, doi: 10.1038/sj.eye.6701788.